


## ILK2/ILK1 Antibody (F40141)


Catalog No.	Formulation	Size
F40141-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F40141-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse
<b>Predicted Reactivity</b>	Rat, Bovine
<b>Format</b>	Purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Purified
<b>UniProt</b>	Q13418
<b>Applications</b>	Western Blot : 1:1000
<b>Limitations</b>	This ILK2/ILK1 antibody is available for research use only.



Western blot analysis of ILK2/ILK1 antibody and HeLa lysate.



Western blot analysis of anti-ILK2/ILK1 antibody and mouse heart tissue lysate

## Description

Transduction of extracellular matrix signals through integrins influences intracellular and extracellular functions, and appears to require interaction of integrin cytoplasmic domains with cellular proteins. Integrin-linked kinase (ILK) is an ankyrin repeat containing 51 kDa receptor-proximate serine-threonine kinase (1), with a reported migration rate of 59K. This 451 amino acid protein interacts with the cytoplasmic domain of the beta-1 integrin subunit and contains sequence motifs found in pleckstrin homology domains capable of interacting with phosphoinositide lipids. ILK is an upstream regulator of  $\text{Pi}(3)\text{K}$  dependant activation of protein kinase B (PKB/AKT) and inhibition of glycogen synthase kinase 3 (GSK-3). ILK2 expression is associated with mediation of cell architecture, adhesion to integrin substrates and anchorage-dependent growth in epithelial cells. ILK2 is overexpressed in some highly invasive tumor cell lines.

## Application Notes

Titration of the ILK2/ILK1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 391-421 from the human protein was used as the immunogen for this ILK2/ILK1 antibody.

## Storage

Aliquot the ILK2/ILK1 antibody and store frozen at  $-20^{\circ}\text{C}$  or colder. Avoid repeated freeze-thaw cycles.