


Interferon alpha 2 Antibody [clone N39] (V2612)

Catalog No.	Formulation	Size
V2612-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2612-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2612SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	N39
Purity	Protein G affinity chromatography
UniProt	P01563
Localization	Cytoplasmic, extracellular (secreted)
Applications	ELISA (order The BSA-free Format For Coating) :
Limitations	This Interferon alpha 2 antibody is available for research use only.



Description

Recognizes a protein of 16-27kDa, identified as human interferon-II) (IFN-(II). Its epitope maps between aa112-148 of IFN-II) (total aa172). This mAb is specific for IFN-(II) and does not cross-react with IFN-(I). The site recognized by this mAb is

called site I and is responsible for the antiviral and anti-proliferative activities of IFN-(II). Epitopes of N27 and N39 mAbs are different and represent a good combination of antibodies to set up an ELISA assay for the quantitation of IFN-(II) after viral infections. The IFN- family consists of 24 or more genes or pseudo-genes. IFN-II) is one of the two distinct families (I and II) of human IFN-. The -interferon are mainly produced by lymphocytes, monocytes, macrophages, and cell lines such as Namalwa and KG1 following induction by viruses, nucleic acids, and glucocorticoid hormones. They are involved in virus resistance on target cells, inhibition of cell proliferation, induction of cytokines and regulation of expression of MHC class I antigens.

Application Notes

Optimal dilution of the Interferon alpha 2 antibody should be determined by the researcher.

Immunogen

Purified recombinant human IFNA2 was used as the immunogen for the Interferon alpha 2 antibody.

Storage

Store the Interferon alpha 2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).