


# Klotho Antibody (Center Region) (F54136)

| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F54136-0.2ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.2 ml  |
| F54136-0.05ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.05 ml |

#### **Bulk quote request**

| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human  |
| Format             | Antigen affinity purified                                |
| Clonality          | Polyclonal (rabbit origin)                               |
| Isotype            | Rabbit Ig  |
| Purity             | Antigen affinity   |
| UniProt            | Q9UEF7   |
| Applications       | Western Blot: 1:2000                                     |
| Limitations        | This Klotho antibody is available for research use only. |



## **Description**

May have weak glycosidase activity towards glucuronylated steroids. However, it lacks essential active site Glu residues at positions 239 and 872, suggesting it may be inactive as a glycosidase in vivo. May be involved in the regulation of calcium and phosphorus homeostasis by inhibiting the synthesis of active vitamin D (By similarity). Essential factor for the specific interaction between FGF23 and FGFR1 (By similarity). [UniProt]

#### **Application Notes**

The stated application concentrations are suggested starting points. Titration of the Klotho antibody may be required due to differences in protocols and secondary/substrate sensitivity.

### **Immunogen**

A portion of amino acids 331-365 from human Klotho was used as the immunogen for the Klotho antibody.

## **Storage**

Aliquot the Klotho antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.