


# **LAMP1 Antibody / CD107A [clone LAMP1/7458] (V4333)**

| Catalog No.    | Formulation   | Size   |
|----------------|---|--------|
| V4333-100UG    | 0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4333-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug  |
| V4333SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                          | 100 ug |

## **Bulk quote request**

| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human  |
| Format             | Purified   |
| Clonality          | Monoclonal (mouse origin)                                  |
| Isotype            | Mouse IgG1, kappa  |
| Clone Name         | LAMP1/7458   |
| Purity             | Protein A/G affinity                                       |
| UniProt            | P11279   |
| Localization       | Cytoplasm, Cell membrane                                   |
| Applications       | Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT |
| Limitations        | This LAMP1 antibody is available for research use only.    |



IHC staining of FFPE human prostate tissue with CD107a / LAMP1 antibody (clone LAMP1/7458). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## **Description**

Lysosome-associated membrane proteins (LAMP) are glycosylated type I membrane proteins that play a role in the biogenesis of the pigment melanin. LAMP-1 (also designated CD107a) and LAMP-2 (also designated CD107b) are involved in a variety of functions, including cellular adhesion, and are thought to participate in the process of tumor invasion and metastasis. Newly synthesized LAMP-1 and LAMP-2 proteins are sorted at the trans-Golgi network and are

transported intracellularly via a pathway that is distinct from the Clathrincoated vesicles used for the mannose-6 phosphate receptor. LAMP-1 is expressed on the surface of Thrombin-activated but not resting platelets, and it is thought to be involved in the adhesive, prothrombic properties of these cells. Both LAMP-1 and LAMP-2 are involved in maintaining lysosome acidity and protecting the lysosomal membranes from autodigestion, and their expression is increased in patients with lysosomal storage disorders.

### **Application Notes**

Optimal dilution of the LAMP1 antibody should be determined by the researcher.

#### **Immunogen**

A recombinant partial protein sequence (within amino acids 200-400) from the human protein was used as the immunogen for the LAMP1 antibody.

#### **Storage**

Aliquot the LAMP1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.