


## LEF1 Antibody (F41886)


| Catalog No.   | Formulation                                | Size    |
|---------------|--|---------|
| F41886-0.4ML  | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml  |
| F41886-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

**Bulk quote request**

|                             |   |
|-----------------------------|---|
| <b>Availability</b>         | 1-3 business days                                       |
| <b>Species Reactivity</b>   | Human   |
| <b>Predicted Reactivity</b> | Mouse, Rat  |
| <b>Format</b>               | Antigen affinity purified                               |
| <b>Clonality</b>            | Polyclonal (rabbit origin)                              |
| <b>Isotype</b>              | Rabbit Ig   |
| <b>Purity</b>               | Antigen affinity  |
| <b>UniProt</b>              | Q9UJU2  |
| <b>Applications</b>         | Western Blot : 1:1000<br>Immunofluorescence : 1:10-1:50 |
| <b>Limitations</b>          | This LEF1 antibody is available for research use only.  |



Fluorescent confocal image of A431 cell stained with LEF1 antibody at 1:25. LEF1 immunoreactivity is localized to the nucleus strongly and cytoplasm weakly.



LEF1 antibody western blot analysis in MDA-MB435 lysate.

## Description

Lymphoid enhancer-binding factor 1 participates in the Wnt signaling pathway. Activates transcription of target genes in the presence of CTNNB1 and EP300. May play a role in hair cell differentiation and follicle morphogenesis. TLE1, TLE2, TLE3 and TLE4 repress transactivation mediated by LEF1 and CTNNB1. Regulates T-cell receptor alpha enhancer function. Binds DNA in a sequence-specific manner. PIAG antagonizes both Wnt-dependent and Wnt-independent activation by LEF1 (By similarity). Isoform 3 lacks the CTNNB1 interaction domain and may be an antagonist for Wnt signaling. Isoform 5 transcriptionally activates the fibronectin promoter, binds to and represses transcription from the E-cadherin promoter in a CTNNB1-independent manner, and is involved in reducing cellular aggregation and increasing cell migration of pancreatic cancer cells. Isoform 1 transcriptionally activates MYC and CCND1 expression and enhances proliferation of pancreatic tumor cells. [UniProt]

## Application Notes

Titration of the LEF1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 10-37 from the human protein was used as the immunogen for this LEF1 antibody.

## Storage

Aliquot the LEF1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.