


LRP1B Antibody (RQ6493)


Catalog No.	Formulation	Size
RQ6493	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q9NZR2
Applications	Immunohistochemistry (FFPE) : 2-5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This LRP1B antibody is available for research use only.




IHC staining of FFPE human ovarian cancer tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human pancreatic cancer tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human liver cancer tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placental tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human lung cancer tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE mouse brain tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat brain tissue with LRP1B antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

Low-density lipoprotein receptor-related protein 1B is a protein that in humans is encoded by the LRP1B gene. This gene encodes a member of the low density lipoprotein (LDL) receptor family. These receptors play a wide variety of roles in normal cell function and development due to their interactions with multiple ligands. Disruption of this gene has been reported in several types of cancer.

Application Notes

Optimal dilution of the LRP1B antibody should be determined by the researcher.

Immunogen

An E. coli-derived human protein (amino acids D3704-K4547) was used as the immunogen for the LRP1B antibody.

Storage

After reconstitution, the LRP1B antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.