


# M6PR Antibody / IGF2R [clone 6G2] (RQ6533)


| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| RQ6533      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

## **Bulk quote request**


| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human  |
| Format             | Antigen affinity purified                                      |
| Clonality          | Monoclonal (mouse origin)                                      |
| Isotype            | Mouse IgG2b  |
| Clone Name         | 6G2  |
| Purity             | Affinity purified  |
| Buffer             | Lyophilized from 1X PBS with 2% Trehalose                      |
| UniProt            | P11717   |
| Localization       | Cytoplasmic  |
| Applications       | Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml |
| Limitations        | This M6PR antibody is available for research use only.         |




IHC staining of FFPE human gastric cancer tissue with M6PR antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liver cancer tissue with M6PR antibody (right) and PBS (left, negative control). HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human placental tissue with M6PR antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) HeLa, 2) U-87 MG and 3) HepG2 cell lysate with M6PR antibody. Predicted molecular weight ~274 kDa.

#### **Description**

Insulin-like growth factor 2 receptor, also called IGF2R or I-MPR is a protein that in humans is encoded by the IGF2R gene. This gene is mapped to 6q25.3. This gene encodes a receptor for both insulin-like growth factor 2 and mannose 6-phosphate, although the binding sites for either are located on different segments of the receptor. This receptor functions in the intracellular trafficking of lysosomal enzymes, the activation of transforming growth factor beta, and the degradation of insulin-like growth factor 2. While the related mouse gene shows exclusive expression from the maternal allele, imprinting of the human gene appears to be polymorphic, with only a minority of individuals showing expression from the maternal allele.

#### **Application Notes**

Optimal dilution of the M6PR antibody should be determined by the researcher.

### **Immunogen**

An E. coli-derived human protein (amino acids F424-R529) was used as the immunogen for the M6PR antibody.

#### **Storage**

After reconstitution, the M6PR antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.