


## MCM6 Antibody [clone MCM6/2999] (V7946)


| Catalog No.    | Formulation  | Size   |
|----------------|--|--------|
| V7946-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V7946-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug  |
| V7946SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                             | 100 ug |

**Bulk quote request**


|                           |   |
|---------------------------|---|
| <b>Availability</b>       | 1-3 business days   |
| <b>Species Reactivity</b> | Human   |
| <b>Format</b>             | Purified  |
| <b>Clonality</b>          | Monoclonal (mouse origin)   |
| <b>Isotype</b>            | Mouse IgG2b, kappa  |
| <b>Clone Name</b>         | MCM6/2999   |
| <b>Purity</b>             | Protein G affinity chromatography                                 |
| <b>UniProt</b>            | Q14566  |
| <b>Localization</b>       | Nuclear   |
| <b>Applications</b>       | Immunohistochemistry (FFPE) : 1-2ug/ml<br>Western Blot : 2-4ug/ml |
| <b>Limitations</b>        | This MCM6 antibody is available for research use only.            |



IHC staining of FFPE human tonsil with MCM6 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon with MCM6 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon carcinoma with MCM6 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.


Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using MCM6 antibody (clone MCM6/2999). These results demonstrate the foremost specificity of the MCM6/2999 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free MCM6 antibody as confirmation of integrity and purity.



Western blot testing of human HeLa cell lysate with MCM6 antibody. Expected molecular weight: 92-105 kDa.

## Description

The mini-chromosome maintenance (MCM) family of proteins, including MCM2, MCM3, MCM6 (Cdc21), MCM5 (Cdc46), MCM6 (Mis5) and MCM7 (Cdc47), are regulators of DNA replication that act to ensure replication occurs only once in the cell cycle. Expression of MCM proteins increases during cell growth, peaking at G1/S phase. The MCM proteins each contain an ATP-binding motif, which is predicted to mediate ATP-dependent opening of double-stranded DNA. MCM proteins are regulated by E2F transcription factors, which induce MCM expression, and by protein kinases, which interact with MCM proteins to maintain the post-replicative state of the cell. MCM2/MCM6 complexes function as substrates for

Cdc2/cyclin B in vitro.

## **Application Notes**

Optimal dilution of the MCM6 antibody should be determined by the researcher.

## **Immunogen**

A recombinant human partial protein (amino acids 228-368) was used as the immunogen for this MCM6 antibody.

## **Storage**

Store the MCM6 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).