


# Melanoma gp100 Antibody / PMEL17 [clone PMEL/2039] (V3929)


Catalog No.	Formulation	Size
V3929-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3929-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3929SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	PMEL/2039
Purity	Protein G affinity chromatography
UniProt	P40967
Localization	Cytoplasmic
Applications	ELISA: 2-4ug/ml (order BSA/azide-free format) Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT Western Blot: 1-2ug/ml
Limitations	This Melanoma gp100 antibody is available for research use only.




IHC testing of human skin with Melanoma gp100 antibody (clone PMEL/2039). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.



IHC testing of human melanoma with Melanoma gp100 antibody (clone PMEL/2039). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Melanoma gp100 antibody (clone PMEL/2039). These results demonstrate the foremost specificity of the PMEL/2039 mAb.<BR>Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD&#39;s) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD&#39;s) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free Melanoma gp100 antibody (clone PMEL/2039) as confirmation of integrity and purity.



Western blot testing of human COLO-38 cell lysate with PMEL17 antibody (clone PMEL/2039).

# **Description**

### **Application Notes**

Optimal dilution of the Melanoma gp100 antibody should be determined by the researcher.

# Immunogen

A portion of amino acids 376-502 from the human protein was used as the immunogen for the Melanoma gp100 antibody.

#### **Storage**

Store the Melanoma gp100 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).