


MGMT Antibody (R31679)


Catalog No.	Formulation	Size
R31679	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
Gene ID	4255
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunocytochemistry : 1-2ug/ml Immunofluorescence (FFPE) : 2-4ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This MGMT antibody is available for research use only.




Immunofluorescent staining of FFPE human MCF7 cells with MGMT antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




ICC staining of FFPE human MCF7 cells with MGMT antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.




IHC-P: MGMT antibody testing of human placenta tissue. HIER: steam section in pH6 citrate buffer for 20 min.




Western blot testing of human 1) Jurkat and 2) MDA-MB-453 cell lysate with MGMT antibody. Predicted molecular weight: ~23 kDa.




Immunofluorescent staining of FFPE human MCF7 cells with MGMT antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human ThP-1 cells with MGMT antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= MGMT antibody.



Western blot testing of human 1) Caco-2 and 2) Raji cell lysate with MGMT antibody. Predicted molecular weight: ~23 kDa.



IHC staining of FFPE human colorectal cancer tissue with MGMT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

O6-alkylguanine DNA alkyltransferase (also known as AGT, MGMT or AGAT) is a protein that in humans is encoded by the MGMT gene. This gene is crucial for genome stability. The protein repairs the naturally occurring mutagenic DNA lesion O6-methylguanine back to guanine and prevents mismatch and errors during DNA replication and transcription. It has been found that methylation of the MGMT promoter in the tumor was associated with longer survival.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the MGMT antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Human partial recombinant protein (AA 2-207) was used as the immunogen for this MGMT antibody.

Storage

After reconstitution, the MGMT antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.