


MMS19 Antibody (RQ8332)


Catalog No.	Formulation	Size
RQ8332	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q96T76
Localization	Nuclear, cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells ELISA : 0.1-0.5ug/ml
Limitations	This MMS19 antibody is available for research use only.




IHC staining of FFPE human colorectal adenocarcinoma tissue with MMS19 antibody.
HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human bladder urothelial carcinoma tissue with MMS19 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placental tissue with MMS19 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human HeLa cells with MMS19 antibody (red). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human Caco-2 and 2) mouse brain tissue lysate with MMS19 antibody. Predicted molecular weight ~113 kDa (multiple isoforms).



Flow cytometry testing of fixed and permeabilized human HeLa cells with MMS19 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= MMS19 antibody.

Description

MMS19 nucleotide excision repair protein homolog is a protein that in humans is encoded by the MMS19 gene. MMS19 (MMS19 nucleotide excision repair homolog), also known as MET18, is a 1,030 amino acid nuclear protein containing seven HEAT repeats that belongs to the MET18/MMS19 family. Via its interactions with TFIIH p80 and TFIIH p89 helicases, MMS19 plays a role in nucleotide excision repair (NER) and RNA polymerase II (Pol II) transcription. MMS19 may also function as a transcriptional coactivator of estrogen receptor. While ubiquitously expressed, highest levels of MMS19 have been found in testis. At least five distinct MMS19 protein isoforms exist, which are produced by alternative splicing events.

Application Notes

Optimal dilution of the MMS19 antibody should be determined by the researcher.

Immunogen

An E.coli-derived human recombinant protein (E239-R998) was used as the immunogen for the MMS19 antibody.

Storage

After reconstitution, the MMS19 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.