


MUNC18-1 Antibody / STXBP1 (R32158)


Catalog No.	Formulation	Size
R32158	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	P61764
Localization	Cytoplasmic, membranous
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This MUNC18-1 antibody is available for research use only.




Immunofluorescent staining of FFPE human HeLa cells with MUNC18-1 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




IHC testing of FFPE human glioma tissue with MUNC18-1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.




IHC testing of FFPE mouse brain with MUNC18-1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE rat brain with MUNC18-1 antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



Western blot testing of 1) human HeLa, 2) rat brain, 3) rat C6 and 4) mouse brain lysate with MUNC18-1 antibody. Expected molecular weight ~69 kDa.



Flow cytometry testing of human A549 cells with MUNC18-1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= MUNC18-1 antibody.

Description

Syntaxin-binding protein 1, also known as Munc18-1, is a protein that in humans is encoded by the STXBP1 gene. By fluorescence in situ hybridization, the STXBP1 gene is mapped to chromosome 9q34.1. This gene encodes a syntaxin-binding protein. The encoded protein appears to play a role in release of neurotransmitters via regulation of syntaxin, a transmembrane attachment protein receptor. Mutations in this gene have been associated with infantile epileptic encephalopathy-4. Alternatively spliced transcript variants have been described.

Application Notes

Optimal dilution of the MUNC18-1 antibody should be determined by the researcher.

Immunogen

Amino acids KEYPAVRYSRGEYKDNALLAQLIQDKLDAYKADD of human MUNC18-1 were used as the immunogen for the MUNC18-1 antibody.

Storage

After reconstitution, the MUNC18-1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.