


## Neurogenin 3 Antibody / NGN3 [clone PCRP-NEUROG3-1E10] (V9690)


Catalog No.	Formulation	Size
V9690-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9690-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9690SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b
<b>Clone Name</b>	PCRP-NEUROG3-1E10
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q9Y4Z2
<b>Localization</b>	Nucleus
<b>Applications</b>	ELISA (order BSA-free Format For Coating) : Immunofluorescence : 1-2ug/ml Flow Cytometry : 1-2ug/million cells
<b>Limitations</b>	This Neurogenin 3 antibody is available for research use only.



Immunofluorescent staining of PFA-fixed human HeLa cells using Neurogenin 3 antibody (green, clone PCRP-NEUROG3-1E10) and phalloidin (red).




FACS staining of PFA-fixed human HeLa cells with Neurogenin 3 antibody (blue, clone PCRP-NEUROG3-1E10) and isotype control (red).



SDS-PAGE analysis of purified, BSA-free Neurogenin 3 antibody (clone PCRP-NEUROG3-1E10) as confirmation of integrity and purity.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Neurogenin 3 antibody (clone PCRP-NEUROG3-1E10). These results demonstrate the foremost specificity of the PCRP-NEUROG3-1E10 mAb. Z- and S-score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SDs) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

## Description

This antigen is a transcription factor that interacts with NKX2-2 to regulation transcription of NEUROD1. The neurogenin family of proteins belongs to the basic helix-loop-helix (bHLH) superfamily and consists of Neurogenin 1, Neurogenin 2 and Neurogenin 3 (also designated ngn3). bHLH members are transcriptional regulators that determine cell fate. Neurogenin 3 is expressed in discrete regions of developing neurons and in the embryonic pancreatic islets of Langerhans. HNF-6 (hepatocyte nuclear factor 6) acts as a positive regulator of Neurogenin 3 by binding to and stimulating the neurogenin gene promoter. Neurogenin 3 is involved in the initial differentiation of the four islets cell types, while a network of transcription factors, including Hlxb9, Isl1, NeuroD, Nkx-2.2, Nkx-6.4, Pax-4, Pax-6, PDX-1 and Mash1, are required for final differentiation. Neurogenin 3 acts upstream of NeuroD in a bHLH cascade. Neurogenin 3 activates the expression of NeuroD at the onset of islet cell differentiation through box sequences E1 and E3 in the NeuroD promoter.

## Application Notes

Optimal dilution of the Neurogenin 3 antibody should be determined by the researcher.

## Immunogen

Recombinant human full-length protein was used as the immunogen for the Neurogenin 3 antibody.

## Storage

Aliquot the Neurogenin 3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

