


NKX2.2 Antibody (V3314)

Catalog No.	Formulation	Size
V3314-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3314-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3314SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Protein A affinity chromatography
Buffer	1X PBS, pH 7.4
UniProt	O95096
Gene ID	4821
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This NKX2.2 antibody is available for research use only.



IHC testing of FFPE Ewings sarcoma with NKX2.2 antibody. HIER: steam sections in 10mM citrate buffer, pH 6.0, for 10-20 min.

Description

Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion

protein specific to Ewing sarcoma, and was shown to be differentially upregulated in Ewing sarcoma on the basis of array-based gene expression analysis. Antibody to NKX2.2 detects a valuable marker for Ewing sarcoma, with a sensitivity of 93% and a specificity of 89%, and aids in the differential diagnosis of small round cell tumors.

Application Notes

Titration of the NKX2.2 antibody may be required for optimal performance.

Immunogen

A human partial recombinant protein was used as the immunogen for this NKX2.2 antibody.

Storage

Store the NKX2.2 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).