


NMI Antibody / N-myc and STAT interactor (R31753)


Catalog No.	Formulation	Size
R31753	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
Gene ID	9111
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 0.5-1ug/ml Immunofluorescence (FFPE) : 2-4ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This NMI antibody is available for research use only.




Immunofluorescent staining of FFPE human A431 cells with NMI antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




IHC staining of FFPE human breast cancer with NMI antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer with NMI antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placenta with NMI antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placenta with NMI antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon cancer with NMI antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human lung cancer with NMI antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) placenta, 2) K562, 3) Caco-2, 4) A431, 5) PC-3 and 6) 293T lysate with NMI antibody. Predicted molecular weight ~35 kDa.



Flow cytometry testing of human U937 cells with NMI antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= NMI antibody.

Description

NMYC interactor (NMI) is a protein that interacts with NMYC and CMYC (two members of the oncogene Myc family), and other transcription factors containing a Zip, HLH, or HLH-Zip motif. NMI also interacts with all [STATs](#) except [STAT2](#) and augments STAT-mediated transcription in response to cytokines [IL2](#) and [IFN-gamma](#). mRNA has low expression levels in all human fetal and adult tissues tested except brain and has high expression in cancer cell line-myeloid leukemias.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the NMI antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Human partial recombinant protein (AA 2-307) was used as the immunogen for this NMI antibody.

Storage

After reconstitution, the NMI antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.