


NRF1 Antibody / Nuclear Respiratory Factor 1 (R31050)


Catalog No.	Formulation	Size
R31050	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
UniProt	Q16656
Applications	Western Blot : 0.5-1ug/ml IHC (FFPE) : 0.5-1ug/ml Immunocytochemistry : 0.5-1ug/ml
Limitations	This NRF1 antibody is available for research use only.




Western blot testing of human 1) U-2 OS and 2) HeLa cell lysate with NRF1 antibody. Expected molecular weight: isoforms from 45-67 kDa.




IHC-P: NRF1 antibody testing of human breast cancer tissue



IHC-P: NRF1 antibody testing of human intestinal cancer tissue



ICC testing of NRF1 antibody and A549 cells



IHC-P: NRF1 antibody testing of rat intestine tissue

Description

Nuclear Respiratory Factor 1 is also known as Alpha-Pal. Efiok et al.(1994) identified genes containing NRF1-binding sequences and found that these could be classified either as cellular proliferation genes, or as genes regulating the growth-responsive metabolic pathways of energy transduction, translation, and replication. Virbasius and Scarpulla (1994) noted that the nuclear-encoded mitochondrial transcription factor [TFAM](#) contains potential binding sites for NRF1, NRF2(GABPA) and SP1 within the promoter region. With use of binding and electrophoretic mobility shift assays, DNase footprinting, and mutation analysis of recombinant proteins, they demonstrated specific and functional binding of [NRF1](#) and [NRF2](#) to the TFAM promoter region.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the NRF1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

An amino acid sequence from the middle region of human Nuclear Respiratory Factor 1 (QHGRELLYAFEDQQTQ) was used as the immunogen for this NRF1 antibody (100% homologous in human, mouse and rat).

Storage

After reconstitution, the NRF1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

