


NRTN Antibody / Neurturin (FY12276)


Catalog No.	Formulation	Size
FY12276	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml	100 ug

Bulk quote request


Availability	1-2 days
Species Reactivity	Human, Mouse, Rat
Format	Lyophilized
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Immunogen affinity purified
Buffer	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
UniProt	Q99748
Applications	ELISA : 0.1-0.5ug/ml Immunohistochemistry : 2-5ug/ml Western Blot : 0.25-0.5ug/ml
Limitations	This NRTN antibody is available for research use only.




Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Western blot analysis of NRTN using anti-NRTN antibody. Lane 1: human SH-SY5Y whole cell lysates, Lane 2: human 293T whole cell lysates, Lane 3: mouse heart tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NRTN antibody at 0.5 ug/ml overnight at 4oC, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal was developed using enhanced chemiluminescent. A specific band was detected for NRTN at approximately 22 kDa. The expected band size for NRTN is at 22 kDa.




Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of human thyroid papillary carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of human thyroid papillary carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.



Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.



Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.



Immunohistochemical staining of NRTN using anti-NRTN antibody. NRTN was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NRTN antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.

Description

NRTN antibody detects Neurturin, encoded by the NRTN gene on chromosome 19p13.3. NRTN antibody is widely used in neuroscience, developmental biology, and neurodegeneration studies. Neurturin is a secreted growth factor of the glial cell line-derived neurotrophic factor (GDNF) family, which supports survival and maintenance of neurons. It plays roles in peripheral nervous system development, innervation of target tissues, and maintenance of dopaminergic neurons.

Structurally, Neurturin is a ~20 kDa secreted protein with a cystine-knot motif characteristic of the TGF-beta superfamily. It is synthesized as a precursor with a signal peptide and pro-region that is cleaved to yield the mature form. Neurturin binds to the GFRalpha-2 receptor, which complexes with RET tyrosine kinase to initiate downstream signaling cascades, including MAPK and PI3K-Akt pathways.

Functionally, Neurturin promotes survival of parasympathetic, sensory, and dopaminergic neurons. It enhances axonal outgrowth, synaptic maintenance, and neuronal regeneration after injury. Researchers use NRTN antibody to investigate neurotrophic signaling, neuron survival, and therapies for neurodegenerative diseases.

Clinically, Neurturin has been investigated for therapeutic potential in Parkinson's disease, where delivery of recombinant Neurturin aims to protect and restore dopaminergic neurons. Altered NRTN expression is linked to gastrointestinal innervation defects and autonomic dysfunction. Because of its restricted receptor usage, Neurturin has selective effects compared to other GDNF family members. NSJ Bioreagents offers NRTN antibody for studies of neuroprotection, development, and disease models.

Experimentally, NRTN antibody is used in western blotting to detect the ~20 kDa protein, in immunohistochemistry to study expression in neural and peripheral tissues, and in ELISA to quantify secreted protein in conditioned media. Co-immunoprecipitation with NRTN antibody identifies binding partners including GFRalpha-2 and RET.

Application Notes

Optimal dilution of the NRTN antibody should be determined by the researcher.

Immunogen

E.coli-derived human Neurturin/NRTN recombinant protein (Position: M1-V197) was used as the immunogen for the NRTN antibody.

Storage

After reconstitution, the NRTN antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.