


## NSFL1C Antibody (F53996)

Catalog No.	Formulation	Size
F53996-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F53996-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse
<b>Predicted Reactivity</b>	Bovine, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	Q9UNZ2
<b>Applications</b>	Western Blot : 1:2000
<b>Limitations</b>	This NSFL1C antibody is available for research use only.



Western blot testing of NSFL1C antibody at 1:2000 + human lysate 1: 293, 2: A431, 3: NCI-H1299, 4: U-251 MG, and 5: mouse brain lysate. Predicted molecular weight ~41 kDa.

## Description

Reduces the ATPase activity of VCP. Necessary for the fragmentation of Golgi stacks during mitosis and for VCP-mediated reassembly of Golgi stacks after mitosis. May play a role in VCP-mediated formation of transitional endoplasmic reticulum (tER) (By similarity). Inhibits the activity of CTSL (in vitro).

## Application Notes

The stated application concentrations are suggested starting points. Titration of the NSFL1C antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## **Immunogen**

Amino acids 214-248 from the human protein were used as the immunogen for this NSFL1C antibody.

## **Storage**

Aliquot the NSFL1C antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.