


## Nucleoli Marker Antibody [clone NCM102-1] (V7035)


| Catalog No.    | Formulation                                                                                     | Size   |
|----------------|-------------------------------------------------------------------------------------------------|--------|
| V7035-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                      | 100 ug |
| V7035-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide                      | 20 ug  |
| V7035SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                                                  | 100 ug |
| V7035IHC-7ML   | Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only* | 7 ml   |

[Bulk quote request](#)


|                           |                                                                                                                                                                                                        |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b> | Human                                                                                                                                                                                                  |
| <b>Format</b>             | Purified                                                                                                                                                                                               |
| <b>Clonality</b>          | Monoclonal (mouse origin)                                                                                                                                                                              |
| <b>Isotype</b>            | Mouse IgG1, kappa                                                                                                                                                                                      |
| <b>Clone Name</b>         | NCM102-1                                                                                                                                                                                               |
| <b>Purity</b>             | Protein G affinity chromatography                                                                                                                                                                      |
| <b>Gene ID</b>            | Unknown                                                                                                                                                                                                |
| <b>Localization</b>       | Nucleoli                                                                                                                                                                                               |
| <b>Applications</b>       | Flow Cytometry : 1-2ug/10 <sup>6</sup> cells<br>Immunofluorescence : 1-2ug/ml<br>Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT<br>Prediluted IHC Only Format : incubate for 30 min at RT (1) |
| <b>Limitations</b>        | This Nucleoli Marker antibody is available for research use only.                                                                                                                                      |



IHC testing of Nucleoli Marker antibody and FFPE human tonsil tissue. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



Immunofluorescent staining of AF488 conjugated Nucleoli Marker antibody and FFPE human colon carcinoma. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



Intracellular FACS testing of human 293 cells with Nucleoli Marker antibody (green) and isotype control (gray).

## Description

The nucleolus is the largest structure in the nucleus of eukaryotic cells, where it primarily serves as the site of ribosome synthesis and assembly. Nucleoli also have other important functions like assembly of signal recognition particles and playing a role in the cell's response to stress. Nucleoli are made of proteins and RNA and form around specific chromosomal regions. [Wiki]

## Application Notes

Titering of the Nucleoli Marker antibody may be required for optimal performance.

2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

A preparation of myeloid leukemia cell nuclei was used as the immunogen.

## Storage

Store the Nucleoli Marker antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).