


Occludin Antibody / OCLN [clone OCLN/8526R] (V5088)

Catalog No.	Formulation	Size
V5088-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5088-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5088SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	OCLN/8526R
Purity	Protein A/G affinity
UniProt	Q16625
Localization	Cell Surface, Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 2-4ug/ml
Limitations	This Occludin antibody is available for research use only.



Western blot testing of human kidney tissue lysate with Occludin antibody. Predicted molecular weight ~59 kDa.



SDS-PAGE analysis of purified, BSA-free Occludin antibody (clone OCLN/8526R) as confirmation of integrity and purity.



IHC staining of FFPE human kidney tissue with Occludin antibody (clone OCLN/8526R). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Occludin is a tetraspan integral membrane protein in epithelial and endothelial tight junction (TJ) structures that can contain two extracellular loops. The protein exists in a variety of phosphorylated forms. Phosphorylation is involved in regulating both the localization and the function of Occludin. Expression of Occludin is upregulated by polyunsaturated fatty acids, increasing trans-endothelial cell resistance and reducing cellular permeability to large molecules. The level of Occludin varies greatly depending on tissue; in brain tissue, Occludin is highly expressed at cell-cell contact sites. Non-neural tissues show lower expression and discontinuous distribution. Up-regulation of epithelial Occludin may play a role in enhancing paracellular permeability and be related to the damage to the tight junction.

Application Notes

Optimal dilution of the Occludin antibody should be determined by the researcher.

Immunogen

A recombinant human protein fragment (within amino acids 282-415) was used as the immunogen for the Occludin antibody.

Storage

Aliquot the Occludin antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.