


P-Selectin Antibody / CD62P (R31776)


| Catalog No. | Formulation | Size |
|-------------|---|--------|
| R31776 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

Bulk quote request


| Availability | 1-3 business days |
|--------------------|---|
| Species Reactivity | Human |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit IgG |
| Purity | Antigen affinity |
| Buffer | Lyophilized from 1X PBS with 2% Trehalose |
| UniProt | P16109 |
| Localization | Cell membrane |
| Applications | Immunohistochemistry (FFPE) : 2-5ug/ml Western Blot : 0.5-1ug/ml ELISA : 0.1-0.5ug/ml (human protein tested); request BSA free format for coating |
| Limitations | This P-Selectin antibody is available for research use only. |




IHC staining of FFPE human spleen tissue with P-Selectin antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human tonsil tissue with P-Selectin antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human tonsil tissue with P-Selectin antibody, HRP-labeled secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human HEL cell lysate with P-Selectin antibody. Predicted molecular weight ~91 kDa but may be observed at higher molecular weights due to glycosylation.