


P53 binding protein 1 Antibody / 53BP1 (RQ7044)


Catalog No.	Formulation	Size
RQ7044	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q12888
Localization	Nuclear
Applications	Western Blot : 0.5-1 ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells Immunofluorescence : 5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This p53 binding protein 1 antibody is available for research use only.




IHC staining of FFPE human breast cancer tissue with p53 binding protein 1 antibody.
HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human bladder epithelial carcinoma tissue with p53 binding protein 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human metaplasia of squamous cells of the renal pelvis tissue with p53 binding protein 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human ovarian cancer tissue with p53 binding protein 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human lung cancer tissue with p53 binding protein 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human moderately differentiated rectal adenocarcinoma tissue with p53 binding protein 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human placental tissue with p53 binding protein 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human placental tissue with p53 binding protein 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human HeLa cells with p53 binding protein 1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of human U-2 OS cells with p53 binding protein 1 antibody. Predicted molecular weight: ~214 kDa but may be observed at up to ~450 kDa.

Description

TP53BP1 (Tumor Protein p53-Binding Protein 1), also called 53BP1, is a protein that in humans is encoded by the TP53BP1 gene. Iwabuchi et al.(1998) mapped the TP53BP1 gene to 15q15-q21 by FISH. Iwabuchi et al.(1994) showed that TP53BP1 binds to the conformationally sensitive central domain of wildtype p53 but not to mutant p53 in vitro. Immunoblot analysis by Iwabuchi et al.(1998) showed that expression of TP53BP1 or TP53BP2 enhances the transactivation function of p53 and induces the expression of p21(CDKN1A). Wang et al.(2002) used small interfering RNA directed against TP53BP1 in mammalian cells to demonstrate that TP53BP1 is a key transducer of the DNA damage checkpoint signal. TP53BP1 was required for p53 accumulation, G2/M checkpoint arrest, and the intra-S-phase checkpoint in response to ionizing radiation.

Application Notes

Optimal dilution of the p53 binding protein 1 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids H1239-H1972) was used as the immunogen for the p53 binding protein 1 antibody.

Storage

After reconstitution, the p53 binding protein 1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

