


PABPN1 Antibody / Polyadenylate-binding protein 2 [clone AAHI-16] (RQ8944)

Catalog No.	Formulation	Size
RQ8944	Antibody in PBS with 0.02% sodium azide, 50% glycerol and 0.4-0.5mg/ml BSA	100 ul


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-2 weeks
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	AAHI-16
Purity	Affinity purified
UniProt	Q86U42
Localization	Nucleus, Cytoplasm
Applications	Western Blot : 1:500 Immunohistochemistry (FFPE) : 1:50
Limitations	This PABPN1 antibody is available for research use only.



IHC staining of FFPE human lung cancer tissue with PABPN1 antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon cancer tissue with PABPN1 antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse brain tissue with PABPN1 antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat brain tissue with PABPN1 antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat brain tissue with PABPN1 antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human HepG2, 2) human RT-4, 3) human MCF7, 4) human Jurkat, 5) rat C6, 6) rat PC-12, 7) mouse RAW264.7 and 8) mouse Neuro-2a cell lysate with PABPN1 antibody. Predicted molecular weight ~35 kDa but published data indicates it can be observed at ~50 kDa.