


Pan Cytokeratin Antibody Cocktail (Acidic + Basic) [clone MonoPoly/4999R] (V4503)

Catalog No.	Formulation	Size
V4503-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4503-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4503SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MonoPoly/4999R
Purity	Protein A/G affinity
UniProt	Q7Z794, Q01546
Localization	Cytoplasm
Applications	Flow Cytometry : 0.5-1ug/million cells Immunofluorescence : 1-2ug/ml Western Blot : 0.5-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This Pan Cytokeratin antibody is available for research use only.



IHC staining of FFPE human tonsil tissue with Pan Cytokeratin antibody (clone MonoPoly/4999R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human basal cell carcinoma with Pan Cytokeratin antibody (clone MonoPoly/4999R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate carcinoma tissue with Pan Cytokeratin antibody (clone MonoPoly/4999R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Application Notes

Optimal dilution of the Pan Cytokeratin antibody should be determined by the researcher.

Immunogen

Recombinant fragments and/or synthetic peptides of human Cytokeratin proteins were used as the immunogen for the Pan Cytokeratin antibody.

Storage

Aliquot the Pan Cytokeratin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.