


PD-L1 Antibody / B7-H1 / CD274 [clone PDL1/2743] (V7823)


Catalog No.	Formulation	Size
V7823-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7823-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7823SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	PDL1/2743
Purity	Protein G affinity chromatography
UniProt	Q9NZQ7
Localization	Cell surface, cytoplasm
Applications	ELISA : order BSA-free format for coating Flow Cytometry : 1-2ug/ml Immunofluorescence : 1-2ug/ml
Limitations	This PD-L1 antibody is available for research use only.



Flow cytometry testing of human Jurkat cells with PD-L1 antibody (clone PDL1/2743); Red=isotype control, Blue= cells alone, Orange= PD-L1 antibody.




Immunofluorescent staining of human Jurkat cells with PD-L1 antibody (clone PDL1/2743, green) and Reddot nuclear stain (red).

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using PD-L1 antibody (clone PDL1/2743). These results demonstrate the foremost specificity of the PDL1/2743 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free PD-L1 antibody (clone PDL1/2743) as confirmation of integrity and purity.

Description

PD-L1 is a checkpoint regulator in immune cells, it is expressed on immune or non-hematopoietic cells. Expression of the protein is seen during pregnancy where it has a role in suppressing the immune system. PD-L1 induces an inhibitory signal in activated T-cells and promotes T-cell apoptosis. It is overexpressed in a number of different cancers where it is believed to play a significant role in the cancer's ability to evade the immune system.

Application Notes

Optimal dilution of the PD-L1 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 39-191 from the human protein was used as the immunogen for this PD-L1 antibody.

Storage

Store the PD-L1 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

