


PMEL17 Antibody / Melanoma gp100 [clone PMEL/2037] (V3930)


Catalog No.	Formulation	Size
V3930-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3930-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3930SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	PMEL/2037
Purity	Protein G affinity chromatography
UniProt	P40967
Localization	Cytoplasmic
Applications	ELISA : 2-4ug/ml (order BSA/azide-free format) Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 1-2ug/ml
Limitations	This PMEL17 antibody is available for research use only.



IHC testing of human melanoma with PMEL17 antibody (clone PMEL/2037). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.




IHC testing of human skin with PMEL17 antibody (clone PMEL/2037). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.


Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using PMEL17 antibody (clone PMEL/2037). These results demonstrate the foremost specificity of the PMEL/2037 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free PMEL17 antibody (clone PMEL/2037) as confirmation of integrity and purity.



Western blot testing of human COLO-38 cell lysate with PMEL17 antibody (clone PMEL/2037).

Description

Application Notes

Optimal dilution of the PMEL17 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 376-502 from the human protein was used as the immunogen for the PMEL17 antibody.

Storage

Store the PMEL17 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

