


PRPF18 Antibody / Pre-mRNA-splicing factor 18 (RQ8001)


Catalog No.	Formulation	Size
RQ8001	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q99633
Localization	Nuclear
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This PRPF18 antibody is available for research use only.




IHC staining of FFPE mouse testis tissue with PRPF18 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat testis tissue with PRPF18 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human urothelial carcinoma tissue with PRPF18 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human ovary serous carcinoma tissue with PRPF18 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




Immunofluorescent staining of FFPE human A549 cells with PRPF18 antibody. HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human SH-SY5Y, 2) rat brain, 3) rat PC-12 and 4) mouse brain tissue lysate with PRPF18 antibody. Predicted molecular weight ~40 kDa.



Flow cytometry testing of fixed and permeabilized human ThP-1 cells with PRPF18 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= PRPF18 antibody.



Flow cytometry testing of fixed and permeabilized human JK-1 cells with PRPF18 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= PRPF18 antibody.

Description

Pre-mRNA splicing occurs in 2 sequential transesterification steps. The protein encoded by this gene is found to be essential for the catalytic step II in pre-mRNA splicing process. It is found in the spliceosome, and contains seven WD repeats, which function in protein-protein interactions. This protein has a sequence similarity to the yeast splicing factor Prp18.

Application Notes

Optimal dilution of the PRPF18 antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids E93-L342) was used as the immunogen for the PRPF18 antibody.

Storage

After reconstitution, the PRPF18 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.