


Recombinant B2M Antibody / Beta-2 Microglobulin [clone B2M/7013R] (V9454)

Catalog No.	Formulation	Size
V9454-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9454-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9454SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	B2M/7013R
Purity	Protein A affinity
UniProt	P61769
Localization	Cell Surface, Extracellular (Secreted)
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant B2M antibody is available for research use only.



IHC staining of FFPE human skin with recombinant B2M antibody (clone B2M/7013R).
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant B2M antibody (clone B2M/7013R) as confirmation of integrity and purity.

Description

Recognizes a protein of 12kDa, identified as beta-2 microglobulin. Major histocompatibility complex (MHC) class 1 molecules bind to antigens for presentation on the surface of cells. The proteasome is responsible for producing these antigens from the components of foreign pathogens. MHC class 1 molecules consist of an alpha heavy chain that contains three subdomains (alpha1, alpha2, alpha3) and a non-covalent associating light chain, known as beta-2-Microglobulin. Beta-2-Microglobulin associates with the alpha3 subdomain of the alpha heavy chain and forms an immunoglobulin domain-like structure that mediates proper folding and expression of MHC class 1 molecules. The alpha1 and alpha2 domains of the alpha heavy chain form the peptide antigen-binding cleft. Mutations in the beta-2-Microglobulin gene can enhance the progression of malignant melanoma phenotypes.

Application Notes

Optimal dilution of the recombinant B2M antibody should be determined by the researcher.

Immunogen

Recombinant full-length human B2M protein was used as the immunogen for the recombinant B2M antibody.

Storage

Aliquot the recombinant B2M antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.