


## Recombinant Basic Cytokeratin Antibody (HMW / Type II) [clone rKRTH/6617] (V9305)

Catalog No.	Formulation	Size
V9305-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9305-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9305SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


### Recombinant MOUSE MONOCLONAL

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rKRTH/6617
Purity	Protein A/G affinity
UniProt	Q01546
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant Basic Cytokeratin antibody is available for research use only.



IHC staining of FFPE human skin with recombinant Basic Cytokeratin antibody (clone rKRTH/6617). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human sebaceous gland tissue with recombinant Basic Cytokeratin antibody (clone rKRT/6617). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Basic Cytokeratin antibody (clone rKRT/6617) as confirmation of integrity and purity.

## Description

The keratins are the typical intermediate filament proteins of epithelia, showing an outstanding degree of molecular diversity. Heteropolymeric filaments are formed by pairing of type I and type II molecules. In humans 54 functional keratin genes exist. They are expressed in highly specific patterns related to the epithelial type and stage of cellular differentiation. This antibody can detect high molecular weight CK1, CK2, CK3, CK4, CK5, CK6, CK7 and CK8. It is usually used in a cytokeratin cocktail with type I cytokeratins antibody.

## Application Notes

Optimal dilution of the recombinant Basic Cytokeratin antibody should be determined by the researcher.

## Immunogen

A synthetic peptide sequence common to Type II Cytokeratins was used as the immunogen for the recombinant Basic Cytokeratin antibody.

## Storage

Aliquot the recombinant Basic Cytokeratin antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.