


## Recombinant Beta-2 Microglobulin Antibody [clone rB2M/7279] (V9431)

Catalog No.	Formulation	Size
V9431-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9431-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9431SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Recombinant Mouse Monoclonal
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	rB2M/7279
<b>Purity</b>	Protein A affinity
<b>UniProt</b>	P61769
<b>Localization</b>	Cell Surface, Extracellular (Secreted)
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This recombinant Beta-2 Microglobulin antibody is available for research use only.



IHC staining of FFPE human prostate tissue with recombinant Beta-2 Microglobulin antibody (clone rB2M/7279). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate tissue with recombinant Beta-2 Microglobulin antibody (clone rB2M/7279). Negative control inset: PBS instead of primary antibody to control for secondary binding. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## Description

Recognizes a protein of 12kDa, identified as beta-2 microglobulin. Major histocompatibility complex (MHC) class 1 molecules bind to antigens for presentation on the surface of cells. The proteasome is responsible for producing these antigens from the components of foreign pathogens. MHC class 1 molecules consist of an alpha heavy chain that contains three subdomains (alpha1, alpha2, alpha3) and a non-covalent associating light chain, known as beta-2-Microglobulin. Beta-2-Microglobulin associates with the alpha3 subdomain of the alpha heavy chain and forms an immunoglobulin domain-like structure that mediates proper folding and expression of MHC class 1 molecules. The alpha1 and alpha2 domains of the alpha heavy chain form the peptide antigen-binding cleft. Mutations in the beta-2-Microglobulin gene can enhance the progression of malignant melanoma phenotypes.

## Application Notes

Optimal dilution of the recombinant Beta-2 Microglobulin antibody should be determined by the researcher.

## Immunogen

Recombinant full-length human B2M protein was used as the immunogen for the recombinant Beta-2 Microglobulin antibody.

## Storage

Aliquot the recombinant Beta-2 Microglobulin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.