


# Recombinant CD30 Antibody / Rabbit Monoclonal [clone Ki-1/1747R] (V7254)

Catalog No.	Formulation	Size
V7254-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7254-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7254SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7254IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


# Recombinant RABBIT MONOCLONAL

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	Ki-1/1747R
Purity	Protein A affinity chromatography
UniProt	P28908
Localization	Cell surface, cytoplasmic
Applications	Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Limitations	This recombinant CD30 antibody is available for research use only.



IHC testing of FFPE human Hodgkin's lymphoma stained with recombinant CD30 antibody (clone Ki-1/1747R). Required HIER: steam sections in pH9, 1mM EDTA for 10-20 min.



## **Description**

CD30 is the receptor for TNFSF8/CD30L. May play a role in the regulation of cellular growth and transformation of activated lymphoblasts. Regulates gene expression through activation of NF-kappa-B. [UniProt]

## **Application Notes**

Optimal dilution of the recombinant CD30 antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

### **Immunogen**

A partial human protein was used as the immunogen for the recombinant CD30 antibody.

### **Storage**

Store the recombinant CD30 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).