

## Recombinant CK8 Antibody / Cytokeratin 8 [clone rB22.1] (V3560)

Catalog No.	Formulation	Size
V3560-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3560-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3560SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3560IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)


<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Recombinant Mouse Monoclonal
<b>Isotype</b>	Mouse IgG1, kappa
<b>Clone Name</b>	rB22.1
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P05787
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT Western Blot : 1-2ug/ml
<b>Limitations</b>	This recombinant CK8 antibody is available for research use only.



Western blot testing of human HCT-116 cell lysate with recombinant CK8 antibody (clone rB22.1).



SDS-PAGE analysis of purified, BSA-free recombinant CK8 antibody (clone rB22.1) as confirmation of integrity and purity.



IHC testing of FFPE human colon tissue with recombinant CK8 antibody (clone rB22.1). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.

## Description

Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). Cytokeratin 8 is primarily found in the non-squamous epithelia and is present in majority of adenocarcinomas and ductal carcinomas. It is absent in squamous cell carcinomas. Hepatocellular carcinomas are defined by the use of antibody that recognizes only cytokeratin 8 and 18. Cytokeratin 8 exists on several types of normal and neoplastic epithelia, including many ductal and glandular epithelia such as colon, stomach, small intestine, trachea, and esophagus as well as in transitional epithelium.

## Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the recombinant CK8 antibody to be titrated up or down for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

A cytoskeletal preparation from human HeLa cells was used as the immunogen for this recombinant CK8 antibody.

## Storage

Store the recombinant CK8 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).