


## Recombinant Galectin 1 Antibody [clone GAL1/2499R] (V7302)

Catalog No.	Formulation	Size
V7302-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7302-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7302SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7302IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG, kappa
<b>Clone Name</b>	GAL1/2499R
<b>Purity</b>	Protein G affinity chromatography
<b>UniProt</b>	P09382
<b>Localization</b>	Cytoplasmic, secreted
<b>Applications</b>	ELISA : 2-4ug/ml (order BSA/azide-free format) Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT Western Blot : 1-2ug/ml
<b>Limitations</b>	This recombinant Galectin 1 antibody is available for research use only.



Western blot testing of human HeLa cell lysate with recombinant Galectin 1 antibody (clone GAL1/2499R). Expected molecular weight ~14 kDa.




Western blot testing of human 1) JEG-3 and 2) K562 cell lysate with recombinant Galectin 1 antibody (clone GAL1/2499R). Expected molecular weight ~14 kDa.




IHC testing of FFPE human prostate carcinoma with recombinant Galectin 1 antibody (clone GAL1/2499R). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min and allowed to cool before testing.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using recombinant Galectin 1 antibody (clone GAL1/2499R). These results demonstrate the foremost specificity of the GAL1/2499R mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD&#39;s) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD&#39;s) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free recombinant Galectin 1 antibody (clone GAL1/2499R) as confirmation of integrity and purity.

## Description

Recombinant Galectin 1 antibody detects galectin-1, a member of the galectin family of beta-galactoside-binding proteins encoded by the LGALS1 gene. Galectin-1 is widely expressed and regulates diverse cellular processes including apoptosis, immune modulation, angiogenesis, and extracellular matrix remodeling. Because of its dual role in normal physiology and disease, Recombinant Galectin 1 antibody is a versatile reagent in immunology, cancer biology, and developmental research.

Galectin-1 is a homodimeric protein of approximately 14 kDa per subunit. Each monomer contains a carbohydrate recognition domain that binds beta-galactosides on glycoproteins and glycolipids. Through these interactions, galectin-1 crosslinks receptors and organizes glycoprotein lattices, influencing signaling, adhesion, and migration. It also binds extracellular matrix components such as laminin and fibronectin, linking it to tissue remodeling.

The Recombinant Galectin 1 antibody clone GAL1/2499R provides consistent and specific detection. Recombinant production ensures batch-to-batch reproducibility, which is critical for long-term studies. Clone GAL1/2499R has been cited in peer-reviewed publications exploring immune tolerance, tumor progression, and neuronal survival. Its

performance in immunohistochemistry, Western blotting, and flow cytometry makes it broadly useful across experimental systems.

Research using clone GAL1/2499R has highlighted galectin-1 as a potent immunoregulatory molecule. It induces apoptosis of activated T cells, promotes regulatory T-cell expansion, and suppresses inflammatory responses. In oncology, galectin-1 expression correlates with tumor angiogenesis, immune evasion, and metastasis. Blocking galectin-1 signaling is being studied as a therapeutic strategy to enhance anti-tumor immunity. Beyond pathology, galectin-1 supports neural development and regeneration, where its trophic effects help maintain neuron survival and plasticity.

NSJ Bioreagents provides this Recombinant Galectin 1 antibody to support research in immunology, oncology, and neuroscience. Alternate terms include LGALS1 antibody, beta-galactoside-binding protein antibody, galaptin antibody, HBL-6 antibody, and lectin galactoside-binding soluble 1 antibody.

## Application Notes

Titration of the recombinant Galectin 1 antibody may be required for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

A human recombinant partial protein corresponding to amino acids 12-108 was used as the immunogen for the recombinant Galectin 1 antibody.

## Storage

Store the recombinant Galectin 1 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).