


Recombinant Human IgG4 Fc Antibody [clone RM217] (R20191)

Catalog No.	Formulation	Size
R20191-100UG	1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	RM217
Purity	Protein A purified from animal origin-free supernatant
UniProt	P01861
Gene ID	3503
Applications	ELISA : 50ng/well-200ng/well (Capture); 0.05-0.2ug/ml (Detection)
Limitations	This recombinant Human IgG4 Fc antibody is available for research use only.




Sandwich ELISA with human plasma using recombinant Human IgG4 Fc antibody as the capture, and [biotinylated anti-human light chains \(<#954;+λ\) antibody RM129](#) as the detect, followed by an AP conjugated streptavidin.



Sandwich ELISA with human IgG4 using recombinant Human IgG4 Fc antibody as the capture, and [biotinylated anti-human light chains \(<#954;+λ\) antibody RM129](#) as the detect, followed by an AP conjugated streptavidin.



Recombinant Human IgG4 Fc antibody specifically reacts to hIgG4. No cross reactivity with IgG1, IgG2, IgG3.



ELISA Titration: the plate was coated with different amounts of hIgG4. A serial dilution of recombinant Human IgG4 Fc antibody was used as the primary and an alkaline phosphatase conjugated anti-rabbit IgG as the secondary.

Description

This recombinant Human IgG4 Fc antibody reacts to the Fc region of hIgG4. No cross reactivity with human IgG1, IgG2, IgG3, IgM, IgA, IgD, IgE, mouse/rat/goat IgG.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant Human IgG4 Fc antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Peptide corresponding to the Fc region of hIgG4 was used as the immunogen for this recombinant Human IgG4 Fc antibody.

Storage

Store the recombinant Human IgG4 Fc antibody at -20oC (with glycerol) or aliquot and store at -20oC (without glycerol).