


Recombinant MART-1 Antibody / Rabbit Monoclonal [clone RMMRT1-1] (V7212)

Catalog No.	Formulation	Size
V7212-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7212-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7212SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7212IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant RABBIT MONOCLONAL

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	RMMRT1-1
Purity	Protein A affinity chromatography
UniProt	Q16655
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Prediluted IHC Only Format : incubate for 30 min at RT (1)
Limitations	This recombinant MART-1 antibody is available for research use only.



IHC testing of formalin-fixed, paraffin-embedded human melanoma stained with recombinant MART-1 antibody (clone RMMRT1-1). HIER: steam sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.



IHC testing of formalin-fixed, paraffin-embedded human melanoma stained with recombinant MART-1 antibody (clone RMMRT1-1). HIER: steam sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min.

Description

MART-1/Melan-A is a protein antigen that is found on the surface of melanocytes. Antibodies against the antigen are used in the medical specialty of anatomic pathology in order to recognize cells of melanocytic differentiation, useful for the diagnosis of a melanoma. The MART-1/Melan-A antigen is specific for the melanocyte lineage, found in normal skin, the retina, and melanocytes, but not in other normal tissues. It is thus useful as a marker for melanocytic tumors (melanomas) with the caveat that it is normally found in benign nevi as well. [Wiki]

Application Notes

The optimal dilution of the recombinant MART-1 antibody for each application should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for this recombinant MART-1 antibody.

Storage

Store the recombinant MART-1 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).