


Recombinant MMP9 Antibody [clone rMMP9/1769] (V3627)

Catalog No.	Formulation	Size
V3627-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3627-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3627SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3627IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rMMP9/1769
Purity	Protein G affinity chromatography
UniProt	P14780
Localization	Cytoplasmic, nuclear, secreted
Applications	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This recombinant MMP9 antibody is available for research use only.



IHC testing of FFPE human spleen with recombinant MMP9 antibody (clone rMMP9/1769). Required HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free recombinant MMP9 antibody (clone rMMP9/1769) as confirmation of integrity and purity.

Description

May play an essential role in local proteolysis of the extracellular matrix and in leukocyte migration. Could play a role in bone osteoclastic resorption. Cleaves KiSS1 at a Gly-|-Leu bond. Cleaves type IV and type V collagen into large C-terminal three quarter fragments and shorter N-terminal one quarter fragments. Degrades fibronectin but not laminin or Pz-peptide.

In the arthritis patient this enzyme might contribute to the pathogenesis of joint destruction and might constitute a useful marker of disease status. [UniProt]

Application Notes

The stated application concentrations are suggested starting points. Titration of the MMP9 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids 603-614 were used as the immunogen for the recombinant MMP9 antibody.

Storage

Store the recombinant MMP9 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).