


Recombinant MPO Antibody / Myeloperoxidase [clone rMPO/6904] (V9329)

Catalog No.	Formulation	Size
V9329-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9329-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9329SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rMPO/6904
Purity	Protein A/G affinity
UniProt	P05164
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 1-2ug/ml
Limitations	This recombinant MPO antibody is available for research use only.




IHC staining of FFPE human bone marrow with recombinant MPO antibody (clone rMPO/6904). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human tonsil tissue with recombinant MPO antibody (clone rMPO/6904). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human spleen tissue with recombinant MPO antibody (clone rMPO/6904). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Western blot testing of human spleen tissue lysate using recombinant MPO antibody (clone rMPO/6904). Expected molecular weight: 59-64 kDa (alpha chain, may be observed at higher molecular weights due to glycosylation), 150+ kDa (glycosylated mature form).



SDS-PAGE analysis of purified, BSA-free MPO antibody (clone rMPO/6904) as confirmation of integrity and purity.

Description

Myeloperoxidase (MPO) also called the peroxidase (POD), is an important marker of bone marrow cells. It is one of the members of the family of heme peroxidase super existing in myeloid cells (mainly neutrophils and monocytes of aniline blue particles). With the deepening of the research on MPO, MPO gene polymorphism has been found to lead to individual for some disease susceptibility differences, with a variety of human development is closely related to the occurrence of diseases. The antibody reacts with neutrophil granulocytes and monocytes in blood and with precursors of granulocytes in the bone marrow. The antibody is useful as an aid for classification of neoplastic tissue, i.e. myeloblasts and immature myeloid cells of acute myelogenous leukemia, progranulocytic leukemia, monomyelocytic leukemia, erythroleukemia and myeloblastoma.

Application Notes

Optimal dilution of the recombinant MPO antibody should be determined by the researcher.

Immunogen

A portion of amino acids 150-250 was used as the immunogen for the recombinant MPO antibody.

Storage

Aliquot the recombinant MPO antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.