


# Recombinant TIMP2 Antibody [clone rTIMP2/2335] (V7896)

Catalog No.	Formulation	Size
V7896-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7896-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7896SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


### Recombinant MOUSE MONOCLONAL

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rTIMP2/2335
Purity	Protein G affinity chromatography
UniProt	P16035
Localization	Cytoplasmic, cell surface, secreted
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant TIMP2 antibody is available for research use only.



IHC staining of FFPE human placenta with recombinant TIMP2 antibody (clone rTIMP2/2335). Required HIER: requires boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant TIMP2 antibody (clone rTIMP2/2335) as confirmation of integrity and purity.

#### **Description**

It recognizes a protein of 21kDa, identified as tissue inhibitor of metalloproteinases-2 (TIMP-2). It is closely related to TIMP-1 and shows the highest binding affinity to both the latent (pro) and active forms of 72kDa Type IV collagenase (also known as MMP-2 or gelatinase A). It also has affinity for the active form of 92kDa Type IV collagenase (also known as MMP-9 or gelatinase B). TIMP-2 inhibits the proteolytic invasiveness of tumor cells and normal placental trophoblast cells.

#### **Application Notes**

Optimal dilution of the recombinant TIMP2 antibody should be determined by the researcher.

#### **Immunogen**

Amino acids DSGNDIYGNPI (N-terminal region) were used as the immunogen for the recombinant TIMP2 antibody.

#### **Storage**

Store the recombinant TIMP2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).