


Recombinant TPO Antibody / Thyroid Peroxidase [clone rTPO/7248] (V4900)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V4900-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4900-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V4900SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |


Recombinant MOUSE MONOCLONAL

Bulk quote request

| Availability | 1-3 business days |
|--------------------|---|
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Recombinant Mouse Monoclonal |
| Isotype | Mouse IgG1, kappa |
| Clone Name | rTPO/7248 |
| Purity | Protein A/G affinity |
| UniProt | P07202 |
| Localization | Cytoplasm, Cell surface |
| Applications | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Limitations | This recombinant TPO antibody is available for research use only. |



IHC staining of FFPE human thyroid carcinoma tissue with recombinant TPO antibody (clone rTPO/7248). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Description

Thyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.

Application Notes

Optimal dilution of the recombinant TPO antibody should be determined by the researcher.

Immunogen

A recombinant fragment of human protein was used as the immunogen for the recombinant TPO antibody.

Storage

Aliquot the recombinant TPO antibody and store frozen at -200C or colder. Avoid repeated freeze-thaw cycles.