


Recombinant xCT Antibody / SLC7A11 [clone SLC7A11/9136R] (V5627)

Catalog No.	Formulation	Size
V5627-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5627-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5627SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	SLC7A11/9136R
Purity	Protein A/G affinity
UniProt	Q9UPY5
Localization	Cytoplasm, membrane
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant xCT antibody is available for research use only.



IHC staining of FFPE human triple negative breast cancer tissue with recombinant xCT antibody (clone SLC7A11/9136R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant xCT antibody (clone SLC7A11/9136R) as confirmation of integrity and purity.

Description

xCT, also known as SLC7A11 (solute carrier family 7, (cationic amino acid transporter, y⁺ system) member 11) or CCB1, is a 501 amino acid multi-pass membrane protein that belongs to the polyamine-organocation superfamily of amino acid transporters. Existing as a disulfide-linked heterodimer with CD98, xCT functions as a member of a heteromeric Na⁺-independent anionic amino acid transport system that specifically facilitates the exchange of anionic amino acids for anionic forms of cystine and glutamate, thereby mediating the formation of glutathione within the cell. Due to its involvement in amino acid transport, xCT is associated with the pathogenesis of glioma-induced neurodegeneration and brain edema, as well as pancreatic cancer. The gene encoding xCT maps to human chromosome 4, which encodes nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes.

Application Notes

Optimal dilution of the recombinant xCT antibody should be determined by the researcher.

Immunogen

A recombinant fragment of human SLC7A11 protein was used as the immunogen for the recombinant xCT antibody.

Storage

Aliquot the recombinant xCT antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.