


## RETN Antibody / Resistin [clone RETN/4326] (V8900)


Catalog No.	Formulation	Size
V8900-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V8900-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V8900SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	RETN/4326
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q9HD89
<b>Localization</b>	Secreted
<b>Applications</b>	Western Blot : 2-4 ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This RETN antibody is available for research use only.




IHC staining of FFPE human spleen tissue with RETN antibody (clone RETN/4326).  
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human spleen tissue with RETN antibody (clone RETN/4326).  
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Western blot testing of human heart tissue lysate using RETN antibody (clone RETN/4326). Predicted molecular weight ~11 kDa.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using RETN antibody (clone RETN/4326). These results demonstrate the foremost specificity of the RETN/4326 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

## Description

The cysteine-rich, adipose tissue-specific, secretory factor resistin (resistance to Insulin) also known as ADSF, is a secreted hormone that potentially links obesity to diabetes. Resistin is rich in serine and cysteine residues and contains a unique cysteine repeat motif. Resistin and the resistin-like molecules share the characteristic cysteine composition and other signature features. Resistin-like expression is highest in proliferative epithelial cells and is markedly increased in tumors, suggesting a role in intestinal proliferation.

## Application Notes

Optimal dilution of the RETN antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 19-108 was used as the immunogen for the RETN antibody.

## Storage

Aliquot the RETN antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

