


RXRG Antibody / RXR gamma [clone PCRP-RXRG-5G6] (V9729)

Catalog No.	Formulation	Size
V9729-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9729-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9729SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	PCRP-RXRG-5G6
Purity	Protein A/G affinity
UniProt	P48443
Localization	Nucleus
Applications	ELISA (order BSA-free Format For Coating) : Flow Cytometry : 1-2ug/million cells
Limitations	This RXRG antibody is available for research use only.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using RXRG antibody (clone PCRP-RXRG-5G6). These results demonstrate the foremost specificity of the PCRP-RXRG-5G6 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD#39;s) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD#39;s) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



FACS staining of PFA-fixed human HeLa cells with RXRG antibody (blue, clone PCRP-RXRG-5G6) and isotype control (red).

Description

Two families of retinoid receptors, RARs and RXRs, have been identified. Retinoic acid receptors (RARs) include RARa, RARb and RARg, each of which have a high affinity for all trans-retinoic acids and belong to the same class of nuclear transcription factors as thyroid hormone receptors, vitamin D3 receptor and ecdysone receptor. The ligand-binding domains of the RARs are highly conserved and RAR isoforms are expressed in distinct patterns throughout development and in the mature organism. Members of the retinoid X receptor (RXR) family, RXR, are activated by 9-cis-RA, a stereo- and photo-isomer of all trans-RA that is expressed in vivo in both liver and kidney and may represent a widely used hormone. As is true for the RAR subfamily, the RXR receptors are closely related to each other both in their DNA-binding and ligand-binding domains and are encoded by separate genes at distinct chromosomal loci.

Application Notes

Optimal dilution of the RXRG antibody should be determined by the researcher.

Immunogen

Recombinant full-length human protein was used as the immunogen for the RXRG antibody.

Storage

Aliquot the RXRG antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.