


\$100A1 Antibody (F41686)


| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F41686-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F41686-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

Bulk quote request

| Availability | 1-3 business days |
|----------------------|--|
| Species Reactivity | Human |
| Predicted Reactivity | Bovine |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity |
| UniProt | P23297 |
| Applications | Western Blot : 1:1000 IHC (Paraffin) : 1:50-1:100 |
| Limitations | This S100A1 antibody is available for research use only. |



S100A1 antibody western blot analysis in MDA-MB435 lysate. Predicted molecular weight: 12-15 kDa.



S100A1 antibody immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue.

Description

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in stimulation of Ca2+-induced Ca2+ release, inhibition of microtubule assembly, and inhibition of protein kinase C-mediated phosphorylation. Reduced expression of this protein has been implicated in cardiomyopathies.

Application Notes

Titration of the S100A1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 45-75 from the human protein was used as the immunogen for this S100A1 antibody.

Storage

Aliquot the S100A1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.