


SEC14L3 Antibody / TAP2 (RQ6436)


Catalog No.	Formulation	Size
RQ6436	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q9UDX4
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence (FFPE) : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This SEC14L3 antibody is available for research use only.




IHC staining of FFPE human rectal cancer tissue with SEC14L3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human gallbladder adenocarcinoma tissue with SEC14L3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer tissue with SEC14L3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human gastric cancer tissue with SEC14L3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat testis tissue with SEC14L3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human HeLa cells with SEC14L3 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human HEK293, 2) human U-87 MG, 3) human PC-3, 4) rat lung, 5) rat liver, 6) mouse lung and 7) mouse liver tissue lysate with SEC14L3 antibody. Predicted molecular weight: 37-46 kDa (multiple isoforms).



Flow cytometry testing of human U937 cells with SEC14L3 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= SEC14L3 antibody.

Description

The protein encoded by this gene is highly similar to the protein encoded by the *Saccharomyces cerevisiae* SEC14 gene. The SEC14 protein is a phosphatidylinositol transfer protein that is essential for biogenesis of Golgi-derived transport vesicles, and thus is required for the export of yeast secretory proteins from the Golgi complex. The specific function of this protein has not yet been determined. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Application Notes

Optimal dilution of the SEC14L3 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids R4-V400) was used as the immunogen for the SEC14L3 antibody.

Storage

After reconstitution, the SEC14L3 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.