


## SKIP Antibody / SNW1 / NCoA-62 [clone PCRP-SNW1-2A1] (V4354)


Catalog No.	Formulation	Size
V4354-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4354-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4354SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a
<b>Clone Name</b>	PCRP-SNW1-2A1
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q13573
<b>Localization</b>	Nucleus, cytoplasm
<b>Applications</b>	Flow Cytometry : 1-2ug/million cells
<b>Limitations</b>	This SKIP antibody is available for research use only.



Flow cytometry testing of PFA-fixed human HeLa cells with SKIP antibody (clone PCRP-SNW1-2A1) followed by goat anti-mouse IgG-CF488 (blue), Red = unstained cells.



SDS-PAGE analysis of purified, BSA-free SKIP antibody (clone PCRP-SNW1-2A1) as confirmation of integrity and purity.



Analysis of a HuProt(TM) microarray containing more than 19,000 full-length human proteins using SKIP antibody (clone PCRP-SNW1-2A1). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a mAb to its intended target. A mAb is considered to specific to its intended target, if the mAb has an S-score of at least 2.5. For example, if a mAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that mAb to protein X is equal to 29.

## Description

Nuclear receptor coactivator NCOA62/SNW1 (also called Nuclear Protein SkiP, SKIIP, Ski-interacting protein) is a member of the SNW gene family, encodes a coactivator that enhances transcription from some Pol II promoters. This coactivator can bind to the ligand-binding domain of the vitamin D receptor and to retinoid receptors to enhance vitamin D-, retinoic acid-, estrogen-, and glucocorticoid-mediated gene expression. It can also interact with poly(A)-binding protein 2 to directly control the expression of muscle-specific genes at the transcriptional level. Finally, the protein may be involved in oncogenesis since it interacts with a region of SKI oncoproteins that is required for transforming activity.

## Application Notes

Optimal dilution of the SKIP antibody should be determined by the researcher.

## Immunogen

A recombinant fragment of human protein was used as the immunogen for the SKIP antibody.

## Storage

Aliquot the SKIP antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.