


SMAD2 Antibody (F53216)

| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F53216-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F53216-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Mouse |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity |
| UniProt | Q15796 |
| Applications | Western Blot : 1:1000 |
| Limitations | This SMAD2 antibody is available for research use only. |



Western blot testing of SMAD2 antibody at 1:1000 dilution + NIH3T3 lysate; Predicted molecular weight: 52~60 kDa.

Description

SMAD2 is a receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD2/SMAD4 complex, activates transcription. May act as a tumor suppressor in colorectal carcinoma. Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator. [UniProt]

Application Notes

Titration of the SMAD2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

This SMAD2 antibody was produced from a rabbit immunized with a KLH conjugated synthetic peptide between 106-140 amino acids from the N-terminal region of human SMAD2.

Storage

Aliquot the SMAD2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.