


Sorting nexin 1 Antibody / SNX1 (RQ7751)


Catalog No.	Formulation	Size
RQ7751	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q13596
Localization	Cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells Immunohistochemistry (FFPE) : 2-5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This Sorting nexin 1 antibody is available for research use only.




Immunofluorescent staining of FFPE human A549 cells with Sorting nexin 1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



IHC staining of FFPE human thyroid cancer tissue with Sorting nexin 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human ovarian cancer tissue with Sorting nexin 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liver tissue with Sorting nexin 1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) HeLa and 2) HepG2 cell lysate with Sorting nexin 1 antibody. Predicted molecular weight ~59 kDa, commonly observed at 59-74 kDa.



Flow cytometry testing of human U-2 OS cells with Sorting nexin 1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Sorting nexin 1 antibody.

Description

Sorting nexin-1 is a protein that in humans is encoded by the SNX1 gene. This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This endosomal protein regulates the cell-surface expression of epidermal growth factor receptor. This protein also has a role in sorting protease-activated receptor-1 from early endosomes to lysosomes. This protein may form oligomeric complexes with family members. This gene results in three transcript variants encoding distinct isoforms.

Application Notes

Optimal dilution of the Sorting nexin 1 antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids Q83-Q505) was used as the immunogen for the Sorting nexin 1 antibody.

Storage

After reconstitution, the Sorting nexin 1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.