


Sorting nexin 4 Antibody / SNX4 (RQ7743)


Catalog No.	Formulation	Size
RQ7743	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	O95219
Localization	Cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This Sorting nexin 4 antibody is available for research use only.




IHC staining of FFPE human thyroid papillary carcinoma tissue with Sorting nexin 4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon adenocarcinoma tissue with Sorting nexin 4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liver cancer tissue with Sorting nexin 4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human glioblastoma tissue with Sorting nexin 4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human 293T, 2) human HeLa, 3) rat PC-12, 4) mouse lung and 5) mouse NIH 3T3 cell lysate with Sorting nexin 4 antibody. Predicted molecular weight ~52 kDa.

Description

Sorting nexin-4 is a protein that in humans is encoded by the SNX4 gene. This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This protein associated with the long isoform of the leptin receptor and with receptor tyrosine kinases for platelet-derived growth factor, insulin, and epidermal growth factor in cell cultures, but its function is unknown. This protein may form oligomeric complexes with family members. Two transcript variants, one protein coding and the other non-protein coding, have been found for this gene.

Application Notes

Optimal dilution of the Sorting nexin 4 antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids M1-M450) was used as the immunogen for the Sorting nexin 4 antibody.

Storage

After reconstitution, the Sorting nexin 4 antibody can be stored for up to one month at 4°C. For long-term, aliquot and

store at -20oC. Avoid repeated freezing and thawing.