


SUR1 Antibody / ABCC8 (RQ4054)


Catalog No.	Formulation	Size
RQ4054	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	Q09428
Applications	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/10^6 cells
Limitations	This SUR1 antibody is available for research use only.



Western blot testing of 1) human placenta, 2) rat brain and 3) mouse brain lysate with SUR1 antibody at 0.5ug/ml. Predicted molecular weight ~177 kDa.



Flow cytometry testing of human A431 cells with SUR1 antibody at 1ug/10^6 cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue=SUR1 antibody.

Description

ATP-binding cassette transporter sub-family C member 8 is a protein that in humans is encoded by the ABCC8 gene. The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug resistance. This protein functions as a modulator of ATP-sensitive potassium channels and insulin release. Mutations and deficiencies in this protein have been observed in patients with hyperinsulinemic hypoglycemia of infancy, an autosomal recessive disorder of unregulated and high insulin secretion. Mutations have also been associated with non-insulin-dependent diabetes mellitus type II, an autosomal dominant disease of defective insulin secretion. Alternatively spliced transcript variants have been found for this gene.

Application Notes

Optimal dilution of the SUR1 antibody should be determined by the researcher.

Immunogen

Amino acids TIQREGTLKDFQRSECQLFEHWKTLMNRQDQELEKETVTERKA from the human protein were used as the immunogen for the SUR1 antibody.

Storage

After reconstitution, the SUR1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.