


TCL1 Antibody [clone TLLP1-1] (V3936)

Catalog No.	Formulation	Size
V3936-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3936-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3936SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	TLLP1-1
Purity	Protein G affinity chromatography
UniProt	P56279
Localization	Nuclear, cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This TCL1 antibody is available for research use only.



IHC testing of FFPE human lymph node with TCL1 antibody (clone TLLP1-1). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.

Description

Enhances the phosphorylation and activation of AKT1, AKT2 and AKT3. Promotes nuclear translocation of AKT1. Enhances cell proliferation, stabilizes mitochondrial membrane potential and promotes cell survival. [UniProt]

Application Notes

The stated application concentrations are suggested starting points. Titration of the TCL1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 2-109 from the human protein was used as the immunogen for the TCL1 antibody.

Storage

Store the TCL1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).